

Paul Yates

- Yates PA**, Burman R, Simpson J, Ponomoreva ON, Thayer MJ, Turker MS. Silencing of mouse Aprt is a gradual process in differentiated cells. *Mol Cell Biol.* 2003 Jul;23(13):4461-70.
- McLaughlin T, Hindges R, **Yates PA**, O'Leary DD. Bifunctional action of ephrin-B1 as a repellent and attractant to control bidirectional branch extension in dorsal-ventral retinotopic mapping. *Development.* 2003 Jun;130(11):2407-18.
- McLaughlin T, Hindges R, **Yates PA**, O'Leary DD. Bifunctional action of ephrin-B1 as a repellent and attractant to control bidirectional branch extension in dorsal-ventral retinotopic mapping. *Development.* 2003 Jun;130(11):2407-18.
- Yates PA**, Roskies AL, McLaughlin T, O'Leary DD. Topographic-specific axon branching controlled by ephrin-As is the critical event in retinotectal map development. *J Neurosci.* 2001 Nov 1;21(21):8548-63.
- Erkman L, **Yates PA**, McLaughlin T, McEvilly RJ, Whisenhunt T, O'Connell SM, Krones AI, Kirby MA, Rapaport DH, Bermingham JR, O'Leary DD, Rosenfeld MG. A POU domain transcription factor-dependent program regulates axon pathfinding in the vertebrate visual system. *Neuron.* 2000 Dec;28(3):779-92.
- Brown A, **Yates PA**, Burrola P, Ortuno D, Vaidya A, Jessell TM, Pfaff SL, O'Leary DD, Lemke G. Topographic mapping from the retina to the midbrain is controlled by relative but not absolute levels of EphA receptor signaling. *Cell.* 2000 Jul 7;102(1):77-88.
- O'Leary DD, **Yates PA**, McLaughlin T. Molecular development of sensory maps: representing sights and smells in the brain. *Cell.* 1999 Jan 22;96(2):255-69.
- Frisen J, **Yates PA**, McLaughlin T, Friedman GC, O'Leary DD, Barbacid M. Ephrin-A5 (AL-1/RAGS) is essential for proper retinal axon guidance and topographic mapping in the mammalian visual system. *Neuron.* 1998 Feb;20(2):235-43.